

Absolute and Delay-Dependent Stability of Equations with a Distributed Delay: a Bridge from Nonlinear Differential to Difference Equations

Elena Braverman^{1,2} and Sergey Zhukovskiy¹

Department of Mathematics and Statistics, University of Calgary,
2500 University Drive N.W., Calgary, AB, Canada T2N 1N4

Abstract

We study delay-independent stability in nonlinear models with a distributed delay which have a positive equilibrium. Such models frequently occur in population dynamics and other applications. In particular, we construct a relevant difference equation such that its stability implies stability of the equation with a distributed delay and a finite memory. This result is, generally speaking, incorrect for systems with infinite memory. If the relevant difference equation is unstable, we describe the general delay-independent attracting set and also demonstrate that the equation with a distributed delay is stable for small enough delays.

AMS Subject Classification: 34K20, 92D25, 34K60, 34K23

Keywords: equations with a distributed delay, global attractivity, permanent solutions, Nicholson's blowflies equation, Mackey-Glass equation.

1 Introduction

In models of population dynamics which are described by an autonomous differential equation

$$\frac{dN}{dt} = f(N) - g(N), \quad (1)$$

where $f(N)$ and $g(N)$ are reproduction and mortality rates, respectively, $f(N) > 0$, $g(N) > 0$ for $N > 0$ and $f(N) > g(N)$ for $0 < N < K$, $f(N) < g(N)$ for $N > K$ (K is the carrying capacity of the environment), the positive equilibrium K is stable: all positive solutions converge to K and are monotone. It was argued that the observed data usually

¹Partially supported by the NSERC Research Grant

²Corresponding author. E-mail *maelena@math.ucalgary.ca*. Fax (403)-282-5150. Phone (403)-220-3956.

oscillates about the carrying capacity; in order to model this phenomenon, it was suggested to introduce delay in the production term

$$\frac{dN}{dt} = f(N(t - \tau)) - g(N); \quad (2)$$

the latter equation can have oscillatory solutions, and the delay incorporated in the right hand side can be interpreted as maturation, production or digestion effects. It is usually assumed that the mortality rate was proportional to the present population level

$$\frac{dN}{dt} = -\mu N(t) + f(N(t - \tau)), \quad \mu > 0. \quad (3)$$

The global behavior of solutions of (3) has been extensively studied in literature, in particular in the cases of negative and positive feedback (see, for example, [1, 2] and references therein), the chaotic behavior is impossible in the case of the monotone feedback [3]. However in the case when $f(x)$ is a unimodal function, i.e., increases for $x < K$ and decreases for $x > K$, there may be delay induced instability and complex dynamics [4, 5]. For a detailed overview of the literature on the dynamics of (3) see the recent papers [6, 7]. It is demonstrated in [6, 7] that if f is a unimodal function and positive equilibrium K of the equation $x_{n+1} = f(x_n)$ is globally asymptotically stable, then all solutions of (3) tend to K . In particular, if f has a negative Schwarzian derivative, then local stability of the equilibrium of the difference equation implies its global attractivity [8]. To the best of our knowledge, the first delay-independent stability conditions were obtained in [9]. In the present paper we will try to answer the general question: what are intrinsic properties of the reproduction function f which allow us to conclude that any solution of the equation with a finite memory converges to the equilibrium? Here we consider both general delays (including integral terms) and continuous functions f which may have multiple extrema, tend to infinity at infinity etc.

As special cases, (3) includes the Nicholson's blowflies equation [10, 11] and the Mackey-Glass equation [4, 13]. The Nicholson's blowflies equation

$$\dot{x}(t) = -\delta x(t) + px(t - \tau)e^{-ax(t - \tau)} \quad (4)$$

was used in [11] to describe the periodic oscillation in Nicholson's classic experiments [10] with the Australian sheep blowfly, *Lucila cuprina*. Equation (4) with a distributed delay was studied in [12], where comprehensive results were obtained for the case $\delta < p < \delta e$.

The Mackey-Glass equation [4, 13]

$$\dot{x}(t) = \frac{ax(t - \tau)}{1 + x^\gamma(t - \tau)} - bx(t) \quad (5)$$

models white blood cells production. Local and global stability of the positive equilibrium for equation (5) with variable delays was studied in [14, 15, 16, 17, 18, 19]; to the best of our knowledge, there are no publications on (5) with a distributed delay.

To incorporate random environment influence, some authors included noise in (4) and studied attractivity conditions, see, for example, [20]. However, in applied problems not only

the derivative but also the delay value can be perturbed. We assume that the production delay is not a constant τ but some distributed value which leads to the equation

$$\dot{x}(t) = r \left[\int_{-\infty}^t f(x(s)) d_s R(t, s) - x(t) \right], \quad (6)$$

where $\int_{t-a}^{t-b} d_s R(t, s)$ is the probability that at time t the maturation delay in the production function is between b and a , where $0 < b < a$. We will assume that very large delays are improbable, substituting $-\infty$ in the lower bound with $h(t) \leq t$ which tends to infinity as $t \rightarrow \infty$. In the present paper we consider a rather general form of f , which includes unimodal functions, as well as functions with several extrema. The only requirement is that $f(x)$ has the only positive fixed point. The main result claims that if this fixed point is a global attractor for all positive solutions of the difference equation

$$x_{n+1} = f(x_n) \quad (7)$$

then all solutions of (6) with positive initial conditions tend to this fixed point as well. To some extent this establishes a link between stable differential equation (1) and difference equation (7) which can undergo a series of bifurcations and even transition to chaos. If (7) is globally stable, so is (6). If the unique positive equilibrium of (7) is unstable, (6) can be stable or not, depending on the delay.

The paper is organized as follows. In Section 2 we prove that all solutions with positive initial conditions are positive and bounded and establish some estimates for the lower and the upper bounds. Section 3 presents sufficient conditions under which all positive solutions converge to the positive equilibrium. In Section 4 delay-dependent stability is investigated. In particular, it is demonstrated that equations are globally attractive for delays small enough; if (7) is unstable, then we can find such delays that the positive equilibrium of (6) is not a global attractor. In Section 5 these results are applied to equations of population dynamics with a unimodal reproduction function and a distributed delay, in particular, to the Nicholson's blowflies and Mackey-Glass equations; some open problems are presented.

2 Boundedness and Estimates of Solutions

We consider the equation with a distributed delay

$$\dot{x}(t) = r(t) \left[\int_{h(t)}^t f(x(s)) d_s R(t, s) - x(t) \right], \quad t \geq 0, \quad (8)$$

and the initial condition

$$x(t) = \varphi(t), \quad t \leq 0. \quad (9)$$

As special cases, (8) includes

1. The integrodifferential equation

$$\dot{x}(t) = r(t) \left[\int_{h(t)}^t K(t, s) f(x(s)) ds - x(t) \right] \quad (10)$$

corresponding to the absolutely continuous $R(t, \cdot)$ for any t . Here

$$\int_{h(t)}^t K(t, s) ds = 1 \quad \text{for any } t, \quad K(t, s) = \frac{\partial}{\partial s} R(t, s) \geq 0$$

is defined almost everywhere.

2. The equation with several concentrated delays

$$\dot{x}(t) = r(t) \left[\sum_{k=1}^m a_k(t) f(x(h_k(t))) - x(t) \right], \quad (11)$$

with $a_k(t) \geq 0$, $k = 1, \dots, m$, where $\sum_{k=1}^m a_k(t) = 1$ for any t . This corresponds to

$$R(t, s) = \sum_{k=1}^m a_k(t) \chi_{(h_k(t), \infty)}(s), \quad \text{where } \chi_I(t) \text{ is the characteristic function of interval } I.$$

Definition. An absolutely continuous in $[0, \infty)$ function $x : \mathbb{R} \rightarrow \mathbb{R}$ is called a *solution of the problem* (8),(9) if it satisfies equation (8) for almost all $t \in [0, \infty)$ and conditions (9) for $t \leq 0$.

The integral in the right hand side of (8) should exist almost everywhere. In particular, for (10) with a locally integrable kernel, φ can be any Lebesgue measurable essentially bounded function. For (11) φ should be a Borel measurable bounded function. For any distribution R the integral exists if φ is bounded and continuous (here we assume f is continuous). Besides, as is commonly set in population dynamics models, $\varphi(t)$ is nonnegative and the value at the initial point is positive.

Consider (8),(9) under the following assumptions.

- (a1) $f : [0, \infty) \rightarrow [0, \infty)$ is a continuous function satisfying Lipschitz condition $|f(x) - f(y)| \leq L|x - y|$, $x, y \geq 0$, $f(0) = 0$, $f(x) > x$ for $0 < x < K$ and $0 < f(x) < x$ for $x > K$;
- (a2) $h : [0, \infty) \rightarrow \mathbb{R}$, is a Lebesgue measurable function, $h(t) \leq t$, $\lim_{t \rightarrow \infty} h(t) = \infty$;
- (a3) $r(t)$ is a Lebesgue measurable essentially bounded on $[0, \infty)$ function, $r(t) \geq 0$ for any $t \geq 0$, $\int_0^\infty r(s) ds = \infty$;
- (a4) $R(t, \cdot)$ is a left continuous nondecreasing function for any t , $R(\cdot, s)$ is locally integrable for any s , $R(t, s) = 0$, $s \leq h(t)$, $R(t, t^+) = 1$. Here $u(t^+)$ is the right side limit of function u at point t .

(a5) $\varphi : (-\infty, 0] \rightarrow \mathbb{R}$ is a continuous bounded function, $\varphi(t) \geq 0$, $\varphi(0) > 0$.

First, let us justify that the solution of (8),(9) exists and is unique.

Denote by $\mathbf{L}^2([t_0, t_1])$ the space of Lebesgue measurable functions $x(t)$ such that

$$Q = \int_{t_0}^{t_1} (x(t))^2 dt < \infty, \quad \|x\|_{\mathbf{L}^2([c,d])} = \sqrt{Q},$$

by $\mathbf{C}([t_0, t_1])$ the space of continuous in $[t_0, t_1]$ functions with the sup-norm.

We will use the following result from the book of Corduneanu [21] (Theorem 4.5, p. 95). We recall that operator N is *causal* (or *Volterra*) if for any two functions x and y and each t the fact that $x(s) = y(s)$, $s \leq t$, implies $(Nx)(s) = (Ny)(s)$, $s \leq t$.

Lemma 1 [21] *Consider the equation*

$$\dot{y}(t) = (\mathcal{L}y)(t) + (\mathcal{N}y)(t), \quad t \in [t_0, t_1], \quad (12)$$

where $\mathcal{L}$ is a linear bounded causal operator, $N : \mathbf{C}([t_0, t_1]) \rightarrow \mathbf{L}^2([t_0, t_1])$ is a nonlinear causal operator which satisfies

$$\|\mathcal{N}x - \mathcal{N}y\|_{\mathbf{L}^2([t_0, t_1])} \leq \lambda \|x - y\|_{\mathbf{C}([t_0, t_1])} \quad (13)$$

for λ sufficiently small. Then there exists a unique absolutely continuous solution of (12) in $[t_0, t_1]$, with the initial function being equal to zero for $t < t_0$.

Theorem 1 *Suppose (a1)-(a5) hold. Then there exists a unique solution of (8),(9).*

Proof. To reduce (8) to the equation with the zero initial function, for any $t_0 \geq 0$ we can present the integral as a sum of two integrals

$$\dot{x}(t) = -r(t)x(t) + r(t) \int_{t_0}^t f(x(s)) d_s R(t, s) + r(t) \int_{t_0}^t f(\varphi(s)) d_s R(t, s), \quad (14)$$

where

$$x(t) = 0, \quad t < t_0, \quad \varphi(t) = 0, \quad t \geq t_0.$$

Here $t_0 \geq 0$ is arbitrary, so we begin with $t_0 = 0$ and proceed to a neighboring t_1 to prove the existence of a local solution. Then in (12)

$$(\mathcal{L}x)(t) = -r(t)x(t), \quad (\mathcal{N}x)(t) = r(t) \int_{t_0}^t f(x(s)) d_s R(t, s) + F(t),$$

where

$$F(t) = r(t) \int_{t_0}^t f(\varphi(s)) d_s R(t, s), \quad x(t) = 0, \quad t < t_0, \quad \varphi(t) = 0, \quad t \geq t_0,$$

and for any $\lambda > 0$ there is t_1 , such that

$$\begin{aligned}
\|\mathcal{N}x - \mathcal{N}y\|_{\mathbf{L}^2([t_0, t_1])} &\leq |r(t)| \left\| \int_{t_0}^t |f(x(s)) - f(y(s))| d_s R(t, s) \right\|_{\mathbf{L}^2([t_0, t_1])} \\
&\leq L \operatorname{ess\,sup}_{t \geq 0} |r(t)| \max_{s \in [t_0, t_1]} |x(s) - y(s)| \left\| \int_{t_0}^t d_s R(t, s) \right\|_{\mathbf{L}^2([t_0, t_1])} \\
&\leq L \operatorname{ess\,sup}_{t \geq 0} |r(t)| \|x(s) - y(s)\|_{C([t_0, t_1])} |t_0 - t_1| \leq \lambda \|x - y\|_{\mathbf{C}([t_0, t_1])}
\end{aligned}$$

for $|t_0 - t_1| \leq \lambda / (L \operatorname{ess\,sup}_{t \geq 0} |r(t)|)$, where L was defined in (a1), here λ can be chosen small enough. By Lemma 1 this implies existence and uniqueness of a local solution for (8). This solution is either global or there exists t_2 such that either

$$\liminf_{t \rightarrow t_2} x(t) = -\infty \quad (15)$$

or

$$\limsup_{t \rightarrow t_2} x(t) = \infty. \quad (16)$$

The initial value is positive, so as far as $x(t) > 0$, the solution is not less than the solution of the initial value problem $\dot{x} + r(t)x = 0$, $x(0) = x_0 > 0$ which is positive and the former case (15) is impossible. In addition, $\dot{x}(t) < 0$ for any

$$x(t) > K, \quad x(t) \geq \max \left\{ \max_{0 \leq s \leq t} x(s), \sup_{s \leq 0} \varphi(s) \right\},$$

which contradicts (16). Thus there exists a unique global solution, which completes the proof. $\square$

Theorem 2 *Suppose (a1)-(a5) hold. Then the solution of (8),(9) is positive for $t \geq 0$.*

Proof. After the substitution

$$y(t) = x(t) \exp \left\{ \int_0^t r(\zeta) d\zeta \right\}, \quad (17)$$

equation (8) becomes

$$\dot{y}(t) = r(t) \exp \left\{ \int_0^t r(s) ds \right\} \int_{h(t)}^t f \left(y(s) \exp \left\{ - \int_0^s r(\zeta) d\zeta \right\} \right) d_s R(t, s), \quad t \geq 0. \quad (18)$$

Thus $y(0) > 0$ and $\dot{y}(t) \geq 0$ as far as $y(s) \geq 0$, $s \leq t$, consequently, $y(t) > 0$ for any $t \geq 0$. Since the signs of $y(t)$ and $x(t)$ coincide, then $x(t) > 0$ for any $t \geq 0$. $\square$

Definition. The solution $x(t)$ of (8),(9) is *permanent* if there exist A and B , $B \geq A > 0$, such that

$$A \leq x(t) \leq B, \quad t \geq 0.$$

In the following we prove permanence of all solutions of (8) with positive initial conditions; moreover, we establish bounds for solutions.

Theorem 3 Suppose (a1)-(a5) hold. Then a solution of (8),(9) is permanent.

Proof. By Theorem 2 the solution is positive for $t \geq 0$. By (a2) there exists $t_0 > 0$ such that $h(t) > 0$, $t \geq t_0$. Since the solution is a continuous positive function, then we can define

$$x_{\min} = \min_{t \in [0, t_0]} x(t) > 0, \quad x_{\max} = \max_{t \in [0, t_0]} x(t). \quad (19)$$

Without loss of generality we assume $x_{\min} < K$, $x_{\max} > K$; otherwise, we can choose $\min\{x_{\min}, \lambda K\}$, $\max\{x_{\max}, K/\lambda\}$, where $0 < \lambda < 1$, as $x_{\min}$ and $x_{\max}$, respectively. By (a1) the following values are positive

$$M = \max_{x \in [x_{\min}, x_{\max}]} f(x), \quad m = \min_{x \in [x_{\min}, x_{\max}]} f(x). \quad (20)$$

Define

$$B = \max \left\{ M, x_{\max}, \max_{x \in [0, K]} f(x) \right\}, \quad A = \min \left\{ m, x_{\min}, \min_{x \in [K, B]} f(x) \right\}. \quad (21)$$

Since $f(x) > x$, $0 < x \leq A$ and $f(x) < x$, $x \geq B$, then there exists $\delta > 0$ such that $f(x) \geq A$ for $A - \delta \leq x \leq B$ and $f(x) \leq B$ for $A \leq x \leq B + \delta$. Let us demonstrate

$$x(t) \in [A, B], \quad t \geq 0. \quad (22)$$

By the definition of A, B we have $x(t_0) \in [A, B]$. Suppose the contrary: $x(t) > B$ or $x(t) < A$ for some $t > t_0$.

First, let $x(t) > B$ for some $t > t_0$. Then $x(t) = B + \varepsilon$ for some $\varepsilon \leq \delta$. Denote

$$S_1 = \{t > t_0 | x(t) > B + \varepsilon\}, \quad t^* = \inf S_1.$$

Since $x(t_0) \leq B$ then the set

$$S_2 = \{t | t_0 \leq t < t^*, x(t) \leq B\}$$

is nonempty, denote $t_* = \sup S_2$. Then $x(t_*) = B$, $x(t^*) = B + \varepsilon$; we also have $B \leq x(t) \leq B + \delta$ and $f(x) \leq B$ in the interval $[t_*, t^*]$, thus the derivative is nonpositive

$$\dot{x}(t) = r(t) \left[\int_{h(t)}^t f(x(s)) d_s R(t, s) - x(t) \right] \leq r(t) \left[\int_{h(t)}^t B d_s R(t, s) - B \right] = 0,$$

which contradicts the assumption $x(t^*) = B + \varepsilon > B = x(t_*)$.

Similarly, let us assume that $x(t) = A - \varepsilon$ for some $\varepsilon > 0$, $\varepsilon < \delta$ and some $t > t_0$. After introducing

$$S_1 = \{t > t_0 | x(t) < A - \varepsilon\}, \quad t^* = \inf S_1, \quad S_2 = \{t | t_0 \leq t < t^*, x(t) \geq A\}, \quad t_* = \sup S_2,$$

we have $x(t_*) = A$, $x(t^*) = A - \delta$, $A - \delta \leq x(t) \leq A$ and $f(x(t)) \geq A$ for $t \in [t_*, t^*]$, hence

$$\dot{x}(t) = r(t) \left[\int_{h(t)}^t f(x(s)) d_s R(t, s) - x(t) \right] \geq r(t) \left[\int_{h(t)}^t A d_s R(t, s) - A \right] = 0.$$

This contradicts the assumption $x(t^*) < x(t_*)$. Consequently (22) is valid for $t \geq t_0$ and thus for any $t \geq 0$, the bounds are positive, so the solution is permanent, which completes the proof. $\square$

Example 1. The statement of Theorem 3 is not valid if we omit the condition $\lim_{t \rightarrow \infty} h(t) = \infty$. Consider the equation

$$\dot{x}(t) = 5x(h(t))e^{-x(h(t))} - x(t), \quad t \geq 0, \quad h(t) \equiv -1, \quad x(t) = t + 1, \quad t \in [-1, 0], \quad (23)$$

which is equivalent to the initial value problem

$$\dot{x}(t) + x(t) = 0, \quad x(0) = 1, \quad (24)$$

its solution $x(t) = e^{-t}$ tends to zero as $t \rightarrow \infty$ and so is not permanent.

3 Absolute Global stability for Stable Difference Equations

One of the main steps in establishing global stability property is the proof of the fact that all nonoscillatory about the equilibrium solutions tend to this equilibrium (see, for example, [22]). For ordinary differential equations all solutions are nonoscillatory, for retarded equations it depends on the delay. Below we demonstrate that convergence of nonoscillatory solutions to the equilibrium is quite a common property which is valid for any reproduction function with a unique positive equilibrium. It can be interpreted as: “if nonoscillatory, solutions of delay equations behave asymptotically similar to ordinary differential equations”.

Definition. A solution $x(t)$ of (8),(9) is *nonoscillatory about K* if there exists $\tau > 0$ such that either $x(t) > K$ or $x(t) < K$ for all $t \geq \tau$. Otherwise, $x(t)$ *oscillates about K* .

Theorem 4 Suppose (a1)-(a5) hold. Any nonoscillatory about K solution of (8),(9) converges to K .

Proof. First, let $x(t) < K$, $t \geq \tau$. Without loss of generality we can assume $\tau = 0$. By (a2) there exists $t_0 \geq 0$ such that $h(t) \geq 0$ for $t \geq t_0$. Denote A as in (21). By Theorem 3 we obtain that $x(t) \geq A$ for any $t \geq 0$. Since f is continuous and $f(x) > x$ for $x < K$, then $m_0 = \inf_{x \in [A, K]} f(x) > A$. There may be two cases: $m_0 = K$ and $m_0 < K$. In the former case, since $x(t) < K$, we have

$$\dot{x}(t) > r(t) \left[\inf_{x \in [A, K]} f(x) - x(t) \right] > 0, \quad t \geq t_0,$$

as far as $x(t) > A$, thus the solution of the delay differential equation is not less than the solution of $\dot{x}(t) = r(t)[K - x(t)]$, $0 < x(t_0) < K$, which is increasing and by (a3) (the integral of $r(t)$ diverges) tends to K .




Figure 1: For an arbitrary reproduction function with one positive equilibrium K we construct a series of such points that eventually a solution is in $[m_j, K]$, if it does not exceed K and is in $[K, M_j]$ if a solution is not less than K .

Consider the latter case $m_0 < K$. By the definition of m_0 and $f(x) > x$ for $x \in [A, K]$ we have $l_0 = \sup\{x < K | f(x) \leq m_0\} < m_0$. Taking any α , $l_0 < \alpha < m_0$ and assuming $x(t) \leq \alpha$ for any t , we obtain

$$\dot{x}(t) = r(t) \left[\int_{h(t)}^t f(x(s)) \, d_s R(t, s) - x(t) \right] \geq r(t)(m_0 - \alpha) > 0,$$

which leads to a contradiction $x(t) \rightarrow \infty$ as $t \rightarrow \infty$ since $\int_0^\infty r(s) \, ds$ diverges. Thus, $x(t_*) \geq \alpha$ for some t_* ; moreover, since $\dot{x}(t) \geq 0$ as $x(t) \leq m_0$ then $x(t) \geq \alpha$ for any $t \geq t_*$. Let $h(t) \geq t_*$, $t > t^*$ for some t^* . By the definition of l_0 and α we have $\tilde{m} = \inf_{x \in [\alpha, K]} f(x) > m_0$ and as far as $x(t) \leq m_0$ and $h(t) \geq t^*$ the following inequality holds

$$\dot{x}(t) = r(t) \left[\int_{h(t)}^t f(x(s)) \, d_s R(t, s) - x(t) \right] \geq r(t)(\tilde{m} - m_0) > 0.$$

Assuming $x(t) \leq m_0$ for any t we again obtain a contradiction. Thus, there exist μ_1 and $t_1 > t_0$ such that $x(\mu_1) \geq m_0$ and $h(t) \geq \mu_1$ for $t \geq t_1$. Then $x(t) \geq m_0$ for any $t \geq \mu_1$ and $x(h(t)) \geq m_0$, $t \geq t_1$.

Further, let $m_1 = \inf_{x \in [m_0, K]} f(x) < K$, here $m_0 < m_1$ since $f(x) > x$ for $0 < x < K$.

Similarly, we obtain $x(t) \geq m_1$ whenever $t > t_2$, for some $t_2 > t_1$.

We continue this process. It can be finite (for example, in Fig. 1 we have $m_2 = K$, where the process stops and we deduce $x(t) \rightarrow K$ as $t \rightarrow \infty$) or infinite (see the branch $x(t) > K$

of Fig. 1). In the infinite case we have an increasing sequence $\{m_j\}$, $m_{j+1} = \min_{x \in [m_j, K]} f(x)$ which does not exceed K , so this sequence has a limit d . Since $f(x)$ is continuous then $d = \min_{x \in [d, K]} f(x)$. If $d < K$ then $f(x) - d$ should attain its minimum in $[d, K]$ but $f(x) > x \geq d$, so this minimum is positive and the equality $d = \min_{x \in [d, K]} f(x)$ leads to a contradiction.

Further, let $x(t) > K$. Similarly, we define B as in (21) and $M_0 = \max_{x \in [K, B]} f(x) < B$. There may be two cases: $M_0 = K$ and $M_0 > K$. In the former case we obtain $x(t) \rightarrow K$ as $t \rightarrow \infty$. Consider the latter case. By Theorem 3 we have $x(t) \leq B$ for any $x \geq 0$. By the definition of M_0 and $f(x) < x$, $x \in [K, B]$ we have $s_0 = \inf\{x > K | f(x) \geq M_0\} > M_0$. Similar to the case $x(t) < K$ we demonstrate that there exists ν_1 such that $x(\nu_1) \leq M_0$ and $x(t) \leq M_0$ for any $t \geq \nu_1$. Let t_1 be such that $h(t) \geq \nu_1$ for $t \geq t_1$. Further, we define $M_1 = \max_{x \in [K, M_0]} f(x) < M_0$. We continue this process, it can be finite or infinite (Fig. 1 illustrates an infinite process for $x(t) > K$). Similar to the case $x(t) < K$ we obtain $x(t) \rightarrow K$ as $t \rightarrow \infty$, which completes the proof. $\square$

Example 2. Let us note that in the case of infinite delays nonoscillatory solutions do not necessarily tend to the positive equilibrium. For example, the solution of the equation

$$\dot{x}(t) + x(t) = \frac{e}{2} x(0) e^{-x(0)}, \quad x(0) = 2,$$

which is $x(t) = \frac{1}{e} + \left(2 - \frac{1}{e}\right) e^{-t}$, tends to $1/e$ while the positive equilibrium is $1 - \ln 2$, the monotone solution is nonoscillatory.

Thus for any reproduction function with a unique fixed point $f(x) = x$ nonoscillatory solutions tend to the equilibrium; this is not, generally, true for oscillatory solutions.

Example 3. Consider the Nicholson's blowflies equation

$$\dot{x}(t) = -\delta x(t) + p x(t - \tau) e^{-ax(t - \tau)}. \quad (25)$$

Denote

$$\tau_k = \frac{1}{\delta \sqrt{aN^*(aN^* - 2)}} \left[\arcsin \left(\frac{\sqrt{aN^*(aN^* - 2)}}{aN^* - 1} \right) + 2\pi k \right], \quad k = 0, 1, 2, \dots, \quad (26)$$

where $N^* = 1/a \ln(p/\delta)$ is a positive equilibrium. If $p > \delta e^2$ then the positive equilibrium is locally asymptotically stable for $\tau \in [0, \tau_0)$ and is unstable (locally, thus it cannot be globally attractive) for $\tau > \tau_0$, and (25) undergoes a Hopf bifurcation at N^* when $\tau = \tau_k$ [23] for $k = 0, 1, 2, \dots$.

Now we prove that absolute (delay-independent) convergence holds in some special cases.

Lemma 2 Suppose (a1)-(a5) and at least one of the following conditions holds:

1. $f(x) < K$ for any $0 < x < K$.
2. Denote by $x_{\max}([a, K])$, $0 \leq a \leq K$, the greatest point in $[a, K]$ where $\max_{x \in [a, K]} f(x)$ is attained and assume

$$f(x_{\max}([a, K])) > K \text{ for some } a \in (0, K) \Rightarrow \min_{x \in [K, f(x_{\max}([a, K]))]} f(x) > a. \quad (27)$$

Then any solution of (8),(9) converges to K .

Proof. 1. By Theorem 3 any solution is permanent: $A \leq x(t) \leq B$, where $A < K < B$, $t \geq 0$, and for some $t_0 \geq 0$ we have $h(t) \geq 0$, $t \geq t_0$. Denote $M_0 = \max_{x \in [K, B]} f(x)$ and $m_0 = \min_{x \in [A, K]} f(x)$.

If $M_0 \leq K$ then the derivative is negative for any $x(t) > K$ and the solution either eventually does not exceed K or is decreasing for any $t \geq t_0$. In the latter case the solution tends to the equilibrium; if it has a different limit, we obtain that the derivative is less than a negative number, which is a contradiction. In the former case, if there exists $t > 0$ such that $x(t) < K$ (otherwise, we have a nonoscillatory case $x(t) \geq K$ where we have already proved convergence), then $x(t) \leq K$ for any t (assuming $x(t^*) > K$ we obtain that the derivative of $x(t)$ is negative almost everywhere, while the function changes from K to $x(t^*) > K$), which is again a nonoscillatory case, by Theorem 4 solution $x(t)$ converges to K .

Thus, we can consider $M_0 > K$ only. Then we introduce

$$M_1 = \max_{x \in [K, M_0]} f(x) < M_0, \quad m_1 = \min_{x \in [m_0, K]} f(x) > m_0, \quad s_1 = \min_{x \in [K, M_0]} f(x)$$

and similarly define M_j, m_j and s_j , $j = 2, 3, \dots$. There exists τ_1 such that $\min\{m_1, s_1\} \leq x(t) \leq M_1$ for $t \geq \tau_1$ and t_1 such that $h(t) \geq \tau_1$, $t \geq t_1$. Similar to the proof of Theorem 4 we obtain that there exists a sequence $\tau_1 \leq \tau_2 \leq \dots \leq \tau_j \leq \dots$ such that

$$\min\{m_j, s_j\} \leq x(t) \leq M_j \quad \text{for } t \geq \tau_j.$$

We assume that all $M_j > K$, otherwise we have an eventually monotone case. Since for any continuous f all three sequences are monotone (nonincreasing M_j and nondecreasing m_j, s_j) and tend to K (see the end of the proof of Theorem 4), then $\lim_{t \rightarrow \infty} x(t) = K$.

2. Now suppose that (27) holds for any $a \geq 0$. We begin with $A \leq x(t) \leq B$ in Theorem 3, $t \geq 0$, where A, B are defined in (21), $h(t) > 0$, $t \geq t_0$. Denote

$$M_0 = \max_{x \in [A, K]} f(x), \quad m_0 = \min_{x \in [K, M_0]} f(x). \quad (28)$$

The case $M_0 = K$ was considered in Part 1, thus we can restrict ourselves to the case $M_0 > K$, maximum is attained at

$$x_0 = x_{\max}([A, K]),$$

here x_0 is the greatest point where the maximum is attained. Let us demonstrate that there exists μ_0 such that $x_0 \leq x(t) \leq M_0$, $t \geq \mu_0$.

If $x(t) > S_0 = \max\{M_0, \max_{x \in [K, B]} f(x)\} = \max_{x \in [A, B]} f(x) \geq K$ for all t then we have a nonoscillatory case and convergence to K , which is a contradiction. Thus, $x(t_*) \leq S_0$ for some t_* ; assuming there is $t^* > t_*$ such that $x(t_*) > S_0$ we obtain that the value of the function at the end of the segment is higher than at the beginning point, while the derivative is nonpositive. So there exists $\tau_0 \geq t_0$ such that $x(t) \leq S_0$ for $t > \tau_0$. If $S_0 \neq M_0$ then $S_1 = \max_{x \in [K, S_0]} f(x) < S_0$. Similarly, we find τ_1 such that $x(t) \leq S_1$ for t large enough. Since the sequence $S_{n+1} = \max_{x \in [K, S_n]} f(x)$ is nonincreasing and tends to K then there exists τ_1 such that $x(t) \leq M_0$, $t \geq \tau_1$. Further, we will consider $t \geq t_1$, where $h(t) \geq \tau_1$ whenever $t > t_1$, only.

If $x(t) < s_0 = \min\{m_0, \min_{x \in [A, K]} f(x)\} = \min_{x \in [A, M_0]} f(x)$ for all t , where m_0 was defined in (28), then we have a nonoscillatory case and convergence to K , which is a contradiction. As above, we prove that $x(t) \geq s_0$ for $t \geq \nu_0$, here $\nu_0 \geq t_0$. If $s_0 < m_0$, then we construct a sequence $s_{n+1} = \min_{x \in [s_n, K]} f(x)$ which tends to K . If $m_0 < K$ then some $s_j \geq m_0$. There is $\mu_0 \geq t_1$ such that $x(t) \geq m_0$, $t \geq \mu_0$. Consequently, we have found μ_0 such that $m_0 \leq x(t) \leq M_0$, $t \geq \mu_0$. Let $h(t) \geq \mu_0$, $t \geq t_2$.

Now, if $m_0 = K$ then for $t \geq t_2$ the solution increases as far as $x(t) < K$. Thus, either $x(t) < K$ for any $t \geq t_2$ and this monotone solution converges to K , or, if $x(t_*) \geq K$ for some t_* , $x(t) \geq K$, $t \geq t_*$, and again we have a nonoscillatory solution which converges to K .

Denote

$$M_1 = \max_{x \in [m_0, M_0]} f(x), \quad m_1 = \min_{x \in [m_0, M_1]} f(x).$$

Let us assume $M_1 > K$, $m_1 < K$ and demonstrate that there is $\mu_1 \geq t_2$ such that $m_1 \leq x(t) \leq M_1$, $x(t) \geq \mu_1$, where $M_1 = f(x_{\max}([m_0, K]))$. In fact, for any $t \geq t_2$ the solution is nonincreasing as far as $x(t) \geq M_1$, which gives an upper bound. Considering t where the equation refers only to the values where this bound is valid, we obtain that the solution is nondecreasing if $x(t) \leq m_1$, which together with $m_1 < K < M_1$ confirms the statement. We continue the induction process

$$M_{n+1} = \max_{x \in [m_n, M_n]} f(x), \quad m_{n+1} = \min_{x \in [m_n, M_{n+1}]} f(x),$$

where $K \geq m_{n+1} > m_n$, $K \leq M_{n+1} < M_n$. This process can be infinitely continued if all $m_n < K$, $M_n > K$ (otherwise, at certain stage we have a “monotone” case which implies convergence), and there exists μ_j , $j = 0, 1, 2, \dots$ such that $m_j \leq x(t) \leq M_j$, $t \geq \mu_j$. Let us assume that there are infinite sequences $\{m_n\}$ and $\{M_n\}$, both are monotone and bounded. Then there exist limits

$$x = \lim_{j \rightarrow \infty} m_j \leq K \leq \lim_{j \rightarrow \infty} M_j = X.$$

Since by the assumptions of the theorem $f(M_j) > m_j$, then $f(X) \geq x$. If $x = f(X)$ then either $x = X = f(X) = X$ and $x(t)$ converges to K or $\min_{y \in [K, f_{\max}([x, K])]} f(y) \leq x$ which contradicts the assumptions of the theorem.

Denote by $x_{\max, j}$, $x_{\min, j}$ the sequences where minimum m_j and maximum M_j are attained, respectively (we recall that we choose maximal $x_{\max, j}$ and minimal $x_{\min, j}$ if an extremum is attained at several points). Then $f(x_{\max, j}) = M_j$, $f(x_{\min, j}) = m_j$, $m_j \leq x_{\max, j} \leq K$, $K \leq x_{\min, j} \leq m_j$, $m_{j+1} > x_{\max, j}$, $M_{j+1} < x_{\min, j}$, which implies that $x_{\max, j}$ and $x_{\min, j}$ tend to x and X , respectively. Let us prove that $x = f(X)$. Assume the contrary: $f(X) > x + 2\varepsilon > 0$. Since $\lim_{j \rightarrow \infty} x_{\min, j} = X$ and f is continuous then there exist n_0 such that $f(x_{\min, j}) > x + \varepsilon$, $j \geq n_0$, but $f(x_{\min, j}) = m_{j+1}$, so $m_{j+1} > x + \varepsilon$, which contradicts the equality $\lim_{j \rightarrow \infty} m_j = \sup_j m_j = x$. Thus $\lim_{t \rightarrow \infty} x(t) = K$, which completes the proof. $\square$

Now we can prove the main result of the present paper.

Theorem 5 *Suppose (a1)-(a5) hold and all positive solutions of the difference equation*

$$x_{n+1} = f(x_n) \quad (29)$$

tend to K . Then any positive solution of (8),(9) converges to K .

Proof. If either conditions of Part 1 of Lemma 2 or (27) hold then K attracts all positive solutions of (8),(9). Let us assume that (27) does not hold (and we do not have a monotone case as in Part 1 of Lemma 2), which means that for some $a \in (0, K)$ we have

$$\min_{x \in [K, f(a)]} f(x) \leq a, \quad \text{or} \quad K < b \leq f(a), \quad f(b) \leq a, \quad \text{where} \quad b \in [K, f(a)].$$

We can assume $f(b) = a$, otherwise, since $f(b) < a$, $f(K) = K > a$ and f is continuous, then there is $c \in [K, b]$ such that $f(c) = a$. Thus, $b > K$ and $f^2(b) = f(f(b)) = f(a) \geq b$. Since $f(x) < x$ for any $x > K$ then for any $x > M := \max_{x \in [0, K]} f(x)$ we have $f^2(x) < x$, then there is a fixed point of f^2 in the segment $[b, M]$, in addition to a fixed point $x = K$. Since the fixed point K of (29) cannot be a global attractor unless K is the only fixed point of f^2 [24] then K is not a global attractor of (29), which completes the proof. $\square$

Here we have not considered the case when f has no positive equilibria; however for completeness we will consider this case as well.

Theorem 6 *Suppose (a2)-(a5) hold, $f(0) = 0$ and $f(x) < x$ for any $x > 0$. Then any positive solution of (8),(9) converges to zero.*

Proof. Let t_0 be such that $h(t) \geq 0$ for $t \geq t_0$. Denote $M_0 = \sup_{t \in [0, t_0]} x(t)$. Since $f(x) < x$ then $x(t) \leq M_0$ for any $t \geq 0$. The solution is decreasing as far as $x > M_1$, where $M_1 = \max_{x \in [0, M_0]} f(x) < M_0$.




Figure 2: If $f(x) < x$ for any $x > 0$ then all positive solutions tend to zero.

Suppose S_1 is the smallest point not exceeding M_0 where this maximum is attained (see Fig. 2). Since $f(x) < x$, then $S_1 > M_1$. Let us choose l_1 such that $M_1 < l_1 < S_1$; as far as $x(t) \geq l_1$ we have $\dot{x}(t) \leq r(t)(M_1 - l_1)$, thus eventually any solution is less than l_1 . Further, for some $d > 0$ we have $M_1 - f(x) \geq d$ if $0 < x \leq l_1$. Thus the derivative at any point $x(t) > M_1$ does not exceed (we recall that $x(t) \leq l_1$) the negative value of $-dr(t)$. Consequently, there exists t_1 such that $x(t) \leq M_1$ for $t \geq t_1$. By induction, we denote $M_{k+1} = \sup_{x \in [0, M_k]} f(x) < M_k$, $k = 1, 2, \dots$, and prove that there exists $t_{k+1} > t_k$ such that $x(t) \leq M_{k+1}$ for $t > t_{k+1}$. Since $\lim_{n \rightarrow \infty} M_n = 0$ then $\lim_{t \rightarrow \infty} x(t) = 0$, i.e., any positive solution tends to zero. $\square$

4 The Case When Stability Is Delay-Dependent

In this section we consider the case when stability properties of (8) depend on the delay. First, we prove that for small delays all solutions tend to positive equilibrium K . Second, we demonstrate that if (27) does not hold then there exists equation (8) with parameters satisfying (a1)-(a5) such that its attracting set is as close to $[m, M]$ as prescribed, where

$$M = \max_{x \in [0, K]} f(x) > K, \quad m = \min_{x \in [K, M]} f(x) \quad (30)$$

and $\max_{x \in [m, K]} f(x) = M$ as well.

Let us note that the Lipschitz condition in (a1) implies $|f(x) - K| \leq L|x - K|$.

Theorem 7 Suppose (a1)-(a5) hold and

$$\limsup_{t \rightarrow \infty} \int_{h(t)}^t r(s) ds < \frac{1}{L+1}. \quad (31)$$

Then any positive solution of (8),(9) converges to K .

Proof. Without loss of generality we can assume that (31) is satisfied for $t \geq 0$. We can find a positive $\lambda < 1$ such that $\int_{h(t)}^t r(s) ds < \frac{\lambda}{L+1}$ for $t \geq 0$. By Theorem 3 any solution is permanent:

$$K - a_0 \leq x(t) \leq K + b_0, \quad t \geq 0, \quad 0 < a_0 < K, \quad b_0 > 0.$$

Let us denote $\alpha = \max\{a_0, b_0\}$ and prove that there exists $t_1 \geq 0$ such that $x(t) \leq K + \lambda\alpha$ for any $t > t_1$. If solution $x(t)$ is nonoscillatory about K then by Theorem 4 it tends to the equilibrium and thus for $\varepsilon = \lambda\alpha > 0$ there exists t_1 such that $x(t) \leq K + \varepsilon$ for $t \geq t_1$. Now assume that $x(t)$ is oscillating, t_0 is such a point that $h(t) \geq 0$ for $t \geq t_0$ and $x(t_*) = K$, $x(\tau_*) = K$, where $x(t) > K$ for $t \in (t_*, \tau_*)$. A continuous function $x(t)$ attains its maximum in $[t_*, \tau_*]$, let $x(t^*) = \max_{s \in [t_*, \tau_*]} x(s) = M_0$. We claim that $x(t^*) \leq K + \lambda\alpha$. Assume the contrary: $x(t^*) > K + \lambda\alpha$. By (a1) we have $\max_{x \in [K, M_0]} f(x) = M_1 < M_0$. Denote

$$M_2 = \max\{M_1, K + \lambda\alpha\}, \quad \tau = \sup\{t \in [t_*, t^*] | x(t) \leq M_2\}.$$

Then $x(t) > M_2 \geq K + \lambda\alpha$, $t \in [\tau, t^*]$.

Since $K - \alpha \leq x(t) \leq K + \alpha$ for any $t \geq 0$ then by (a1)

$$\begin{aligned} \dot{x}(t) &= r(t) \left[\int_{h(t)}^t f(x(s)) ds R(t, s) - x(t) \right] \leq r(t) \left[\max_{x \in [K-\alpha, K+\alpha], x > 0} f(x) - (K - \alpha) \right] \\ &= r(t) \left[\max_{x \in [K-\alpha, K+\alpha], x > 0} [f(x) - K] + \alpha \right] \leq r(t)[L\alpha + \alpha] = \alpha(L+1)r(t). \end{aligned} \quad (32)$$

We remark that $\dot{x}(t) < 0$ whenever the following conditions hold: $t \in (\tau, t^*)$, $h(t) \geq t_*$ (all referred prehistory of the solution is between K and $M_1 < M_0$) and $x(t) > M_2 \geq M_1$. Since $f(\tau) = M_2$, $f(t^*) > M_2$, then there are points in (τ, t^*) where the derivative is positive, thus $h(t) < t_*$. Let $\bar{t}$ be such a point. As $r(t) \geq 0$ and $[t_*, \tau] \subset [h(\bar{t}), \bar{t}]$, then

$$\int_{t_*}^{\tau} r(s) ds \leq \int_{h(\bar{t})}^{\bar{t}} r(s) ds < \frac{\lambda}{L+1},$$

consequently,

$$x(\tau) - x(t_*) = \int_{t_*}^{\tau} \dot{x}(s) ds \leq \int_{t_*}^{\tau} \alpha(L+1)r(s) ds < \alpha(L+1)\frac{\lambda}{L+1} = \lambda\alpha.$$

Hence $x(\tau) < x(t_*) + \lambda\alpha = K + \lambda\alpha$, which contradicts the assumption $x(\tau) \geq K + \lambda\alpha$. If we denote $t_1 = t_*$ then $x(t) < K + \lambda\alpha$ for any $t \geq t_1$.

Similar to the previous case, we can prove that there exists τ_1 such that $x(t) \geq K - \lambda\alpha$ for $t \geq \tau_1$.

Thus, we accept $K - \lambda\alpha \leq x(t) \leq K + \lambda\alpha$ as new solution bounds and consider τ_2 such that $h(t) \geq \max\{t_1, \tau_1\}$ for $t \geq \tau_2$. We repeat the induction step and obtain that there exists $t_2 \geq \tau_2$ such that

$$K - \lambda^2\alpha \leq x(t) \leq K + \lambda^2\alpha, \quad t \geq t_2,$$

and a sequence of points $t_1 \leq t_2 \leq t_3 \leq t_n \leq \dots$ such that

$$K - \lambda^3\alpha \leq x(t) \leq K + \lambda^3\alpha, \quad t \geq t_3, \dots, K - \lambda^n\alpha \leq x(t) \leq K + \lambda^n\alpha, \quad t \geq t_n, \dots$$

Since $0 < \lambda < 1$ then $\lim_{t \rightarrow \infty} x(t) = K$, which completes the proof. $\square$

Corollary 1 Suppose (a1)-(a2), (a4)-(a5) hold, $r > 0$ and $\sup_{t \geq 0} (t - h(t)) < \frac{1}{r(L+1)}$ (we recall that L is the Lipschitz constant defined in (a1)). Then all solutions of the equation

$$\dot{x}(t) = r \left[\int_{h(t)}^t f(x(s)) d_s R(t, s) - x(t) \right] \quad (33)$$

tend to K .

Remark 1 Let us remark that the Lipschitz constant L in (31) can be substituted by, generally, a smaller value of

$$L^* = \sup_{x \geq 0} \left| \frac{f(x) - K}{x - K} \right|, \quad (34)$$

since in (32) only $|f(x) - K|$ is estimated for $|x - K| < \alpha$.

Next, let us estimate the attracting set when (8) is not absolutely stable. We introduce m, M as in (30) and denote by $x_{\max}$ the greatest point in $[a, K]$ where $M = \max_{x \in [0, K]} f(x)$ is attained, by $x_{\min}$ the minimal point where $m = \min_{x \in [K, M]} f(x)$ is attained.

Theorem 8 Suppose $f(x)$ satisfies (a1) and

$$m = f(x_{\min}) < x_{\max} < K. \quad (35)$$

Then for any $a \in (m, x_{\max})$, and any $b \in (K, M)$ such that $\min_{x \in [K, b]} f(x) < a$ there exists a problem (8), (9) with parameters satisfying (a2)-(a5) such that

$$\liminf_{t \rightarrow \infty} x(t) = a, \quad \limsup_{t \rightarrow \infty} x(t) = b. \quad (36)$$




Figure 3: Assuming a and b as in the figure exist, we can construct a delay equation such that $[a, b]$ is its attracting set.

Proof. Let us fix $a \in (m, x_{\max})$, and $b \in (K, M)$ such that $d = \min_{x \in [K, b]} f(x) < a$, see Fig. 3.

Then there exists x_1 such that $K < x_1 < b$ and $m_1 = f(x_1) < a$. For the initial function

$$\varphi(t) = a + (t+1)(b-a), \quad t \in [-1, 0],$$

we have $\varphi(-1) = a$, $\varphi(0) = b$ and φ changes continuously from a to b . Since $a < K < x_1 < b$, then $x(s_0) = x_1$ for some $s_0 \in (-1, 0)$. We consider the equation

$$\dot{x}(t) = r[f(x(h(t))) - x(t)], \quad t \geq 0, \quad (37)$$

where $r > 0$. Obviously parameters of (37) and the initial function satisfy (a3)-(a5).

We choose $h(t) = s_0$, $t \in [0, \tau_1]$, where $\tau_1 = \frac{1}{r} \ln \left(\frac{b-m}{a-m} \right) > 0$. Then

$$x(t) = (b-m)e^{-rt} + m, \quad t \in [0, \tau_1],$$

so $x(\tau_1) = a$ and $x(t)$ changes continuously from b to a in $[0, \tau_1]$, $a < x_{\max} < b$, thus there exists $s_1 \in (0, \tau_1)$ such that $x(s_1) = x_{\max}$. We assume $h(t) = s_1$, $t \in (\tau_1, \tau_2]$, where $\tau_2 = \tau_1 + \frac{1}{r} \ln \left(\frac{M-a}{M-b} \right) > \tau_1$. The solution is

$$x(t) = (a-M)e^{-r(t-\tau_1)} + M, \quad t \in [\tau_1, \tau_2],$$

hence $x(\tau_2) = b$. We continue periodically with $h(t)$ piecewise constant such that

$$x(h(t)) = \begin{cases} x_1, & t \in (\tau_{2n}, \tau_{2n+1}], \\ x_{\max}, & t \in (\tau_{2n+1}, \tau_{2n+2}], \end{cases} \quad n = 0, 1, 2, \dots,$$

where $\tau_0 = 0$, $h(t) \in (\tau_{k-1}, \tau_k)$ if $t \in (\tau_k, \tau_{k+1})$, $x(\tau_{2n}) = b$, $x(\tau_{2n+1}) = a$, $n = 0, 1, 2, \dots$ and $a \leq x(t) \leq b$ for any t . Here the delay is bounded and piecewise continuous, it obviously satisfies (a2). $\square$

Corollary 2 *If (a1)-(a5) and (35) hold then the sharpest attracting interval for (8),(9) is (m, M) , where m and M are defined in (30).*

Remark 2 *Suppose the difference equation $x_{n+1} = f(x_n)$ has a stable 2-cycle (a, b) in such a way that $f(f(x)) < x$ for $a < x < K$, $x > b$ and $f(f(x)) > x$ for $x < a$, $K < x < b$. Then for any $\varepsilon > 0$ there exist such distributed delay and initial function that $[a + \varepsilon, b - \varepsilon]$ is an attracting interval of (8),(9). Moreover, if $K \in (a, b)$, function f satisfies (a1), $f(a) > b$ and $f(b) < a$, then, similar to the proof of Theorem 8 we can demonstrate that there exists problem (8),(9) such that (a, b) is its attracting interval.*

Remark 3 *The claims of Corollary 2 and Remark 2 to some extent complement Theorem 6 in [7] where the sharpest invariant and attracting interval was found in the case when absolute stability fails but there exists a unique globally attractive 2-cycle for the relevant difference equation (for equations with a unique constant concentrated delay).*

5 Applications and Discussion

The global attractivity results can be applied to any unimodal function f and any type of finite delay. For example, we can deduce the following result in the stream of [6], see also [8] and Proposition 2.1 in [25].

Theorem 9 *Suppose (a1)-(a5) hold, f is three times continuously differentiable and has the only critical point $x_0 > 0$ (maximum),*

$$(Sf)(x) = \frac{f'''(x)}{f'(x)} - \frac{3}{2} \left(\frac{f''(x)}{f'(x)} \right)^2 < 0 \quad \text{for } x \neq x_0$$

and $|f'(K)| \leq 1$. Then any solution of (8),(9) tends to K as $t \rightarrow \infty$.

As an application, consider the Nicholson's blowflies equation with a distributed delay

$$\dot{x}(t) = -\delta x(t) + p \int_{h(t)}^t x(s) e^{-ax(s)} d_s R(t, s), \quad (38)$$

where (a2)-(a5) hold.

If $p < \delta$ then by Theorem 6 all positive solutions of (38) go to extinction, i.e., $x(t) \rightarrow 0$ as $t \rightarrow \infty$. If $1 < \frac{p}{\delta} < e^2$ then by Theorem 5 all positive solutions of (38) tend to the positive equilibrium $K = \frac{1}{a} \ln \left(\frac{p}{\delta} \right)$. However, since sustainable oscillations were observed in experiments [10], we speculate that in the real problem the ratio p/δ was above e^2 and delays significant enough.

Particular cases of (38) are the equation with a variable delay

$$\dot{x}(t) = -\delta x(t) + px(h(t))e^{-ax(h(t))}, \quad (39)$$

including the original equation (4) with a constant delay, the integrodifferential equation

$$\dot{x}(t) = -\delta x(t) + p \int_{h(t)}^t k(t-s)x(s)e^{-ax(s)} ds, \quad (40)$$

where

$$\int_{h(t)}^t k(t-s) ds = 1 \quad \text{for any } t \geq 0, \quad (41)$$

and the mixed equation

$$\dot{x}(t) = -\delta x(t) + \alpha p x(g(t))e^{-ax(g(t))} + (1-\alpha)p \int_{h(t)}^t k(t-s)x(s)e^{-ax(s)} ds, \quad 0 \leq \alpha \leq 1. \quad (42)$$

By Theorems 5, 6 and 7 we conclude the following result.

Theorem 10 *If $p \leq \delta$ then all positive solutions of (38) (and thus any of (39), (40), (42)) go to extinction, if $\delta < p < \delta e^2$, all positive solutions tend to the positive equilibrium. If $p \geq e^2$ and*

$$\limsup_{t \rightarrow \infty} (t - h(t)) < \frac{1}{p + \delta}, \quad (43)$$

then all positive solutions of (38) tend to the positive equilibrium.

Proof. The results for $p < \delta e^2$ are immediate corollaries of Theorems 5 and 6. Since the absolute value of the derivative of $f(x) = \frac{p}{\delta} x e^{-ax}$ does not exceed p/δ (which is attained at $x = 0$), then (31) implies (43). $\square$

Remark 4 *We can apply Remark 1 to improve condition (43). In fact, L^* defined in (34) does not exceed $p/(\delta e^2)$ (the absolute value of the minimum of the derivative which is attained at $x = 2/a$) if $p \geq \delta e^2$ and the stability condition becomes*

$$\limsup_{t \rightarrow \infty} (t - h(t)) < \frac{1}{p/e^2 + \delta}. \quad (44)$$

We also notice that for unimodal functions such that their equilibrium (the fixed point $x = f(x)$) exceeds maximum point $x_{\max}$ and such that the minimum of the derivative is less than -1 (otherwise, under some additional conditions which are satisfied for the Nicholson's blowflies and the Mackey-Glass equations, the relevant difference equations are stable [25]), then L^ defined in (34) does not exceed the absolute value of the minimum of the derivative.*

Really, function $g(x) = \left| \frac{f(x) - K}{x - K} \right|$ satisfies $g(0) = 1$, $0 < g(x) \leq -\min_{x \in [x_{\max}, \infty)} f'(x)$ in $[x_{\max}, \infty)$. The value in the right hand side of the latter inequality exceeds 1 and g is decreasing in $[0, x^]$, where $x^* < x_{\max}$ is such point that $f(x^*) = K$, while in $[x^*, x_{\max})$ function $g(x)$ is less than $g(x_{\max})$, which does not exceed the absolute value of the minimum of the derivative.*

Let us compare Theorem 10 and Remark 4 to some known global stability results for the Nicholson's blowflies equation. For example, [22] contains the following global stability condition

$$(e^{\delta\tau} - 1) \left(\frac{p}{\delta} - 1 \right) < 1 \quad (45)$$

for equation (4) with a constant delay.

Condition (45) is slightly sharper for (4) than (43) but (43) is applicable to equations with variable delays and integrodifferential equations as well. Condition (44) is sharper than (45), see Fig. 4. It is also applicable to equations with a variable and/or distributed delay.




Figure 4: The values of delay for which (4) is asymptotically stable are given for $\delta = 1$, $p \geq \delta e^2 > 7$ and conditions (44), (43) and (45), respectively.

For completeness of references, we remark that oscillation of (38) was studied in [26].

All the main results are also relevant for the Mackey-Glass equation with a distributed delay

$$\dot{x}(t) = \int_{h(t)}^t \frac{ax(s)}{1+x^\gamma(s)} ds R(t,s) - bx(t), \quad a > 0, \quad b > 0, \quad \gamma > 0, \quad (46)$$

which also involves the equation

$$\dot{x}(t) = -bx(t) + \frac{\alpha ax(g(t))}{1+x^\gamma(g(t))} + (1-\alpha)a \int_{h(t)}^t k(t-s) \frac{x(s)}{1+x^\gamma(s)} ds, \quad 0 \leq \alpha \leq 1, \quad (47)$$

where (41) is satisfied, as well as its special cases of the equation with a variable delay ($\alpha = 1$) and the integrodifferential equation ($\alpha = 0$). If $a < b$ then all solutions of (46) go to extinction. If $a > b$ and

$$\gamma < \frac{2a}{a-b} \quad (48)$$

then all solutions tend to the positive equilibrium $K = \left(\frac{a}{b} - 1 \right)^{1/\gamma}$.

Again, Theorems 5, 6 and 7 imply the following result.

Theorem 11 *If $a \leq b$ then all positive solutions of (46) (and thus of (47)) go to extinction. If either $0 < \gamma \leq 2$ or $\gamma > 2$ and $b < a < \frac{\gamma b}{\gamma - 2}$, then all positive solutions tend to the positive equilibrium K . If $\gamma > 2$, $a \geq \frac{\gamma b}{\gamma - 2}$ and*

$$\limsup_{t \rightarrow \infty} (t - h(t)) < \frac{4b^2\gamma}{a^2(\gamma - 1)^2 + 4ab\gamma}, \quad (49)$$

then all positive solutions of (46) tend to the positive equilibrium.

Proof. We remark that for $\gamma \leq 1$ the function $f(x) = \frac{ax}{b(1 + x^\gamma)}$ is monotone increasing, and we have stability for any delay. For $1 \leq \gamma \leq 2$ the relevant difference equation $x_{n+1} = f(x_n)$ is locally asymptotically stable, we apply Theorem 9. It is known that for the relevant difference equation local asymptotic stability implies global asymptotic stability (see [25], Example 3.2), thus the difference equation is globally asymptotically stable whenever

$$|f'(K)| = \left| 1 - \gamma + \frac{b}{a}\gamma \right| < 1, \quad \text{where } \gamma > 2,$$

or $a < \gamma b/(\gamma - 2)$. According to Remark 4, we can take the absolute value of the minimum of the derivative which is attained at $\left(\frac{\gamma + 1}{\gamma - 1}\right)^{1/\gamma}$ as L^* :

$$L^* = \frac{a(\gamma - 1)^2}{4b\gamma}.$$

Application of Theorem 7 and Remark 4 completes the proof. $\square$

Finally, let us formulate some open problems.

1. Generally, absolute stability results are incorrect for systems with infinite memory. Deduce stability results when (a2) is not satisfied but measure $d_s R(t, s)$ decays exponentially with memory

$$\int_{-\infty}^{t-\tau} d_s R(t, s) \leq S e^{-\alpha\tau}.$$

2. For arbitrary $f(x)$ satisfying (a1) and the equation with a constant delay, characterize attractive sets.
3. Obtain stability results for equations with a distributed delay in the multistability case: there are several points satisfying $f(x) = x$. If K_1 is the first positive fixed point and $f(x) < x$ for $0 < x < K_1$, describe initial conditions leading to extinction (Alley effect).

4. Would the results of the present paper remain valid for the equation

$$\dot{x}(t) = r(t) \left[f \left(\int_{h(t)}^t x(s) d_s R(t, s) \right) - x(t) \right], \quad t \geq 0,$$

if (a1)-(a5) hold?

References

- [1] T. Krisztin, H. O. Walther and J. Wu, Shape, smoothness and invariant stratification of an attracting set for delayed monotone positive feedback. Fields Institute Monographs, 11. American Mathematical Society, Providence, RI, 1999.
- [2] T. Krisztin and H. O. Walther, Unique periodic orbits for delayed positive feedback and the global attractor, *J. Dynam. Differential Equations* **13** (2001), no. 1, 1–57.
- [3] J. Mallet-Paret and G. R. Sell, The Poincaré-Bendixson theorem for monotone cyclic feedback systems with delay, *J. Differential Equations* **125** (1996), 441–489.
- [4] M. C. Mackey and L. Glass, Oscillation and chaos in physiological control systems, *Science* **197** (1977), 287–289.
- [5] B. Lani-Wayda, Erratic solutions of simple delay equations, *Trans. Am. Math. Soc.* **351** (1999), 901–945.
- [6] G. Röst and J. Wu, Domain-decomposition method for the global dynamics of delay differential equations with unimodal feedback, *Proc. R. Soc. Lond. Ser. A Math. Phys. Eng. Sci.* **463** (2007), no. 2086, 2655–2669.
- [7] E. Liz and G. Röst, On the global attractor of delay differential equations with unimodal feedback, submitted.
- [8] D. Singer, Stable orbits and bifurcation of maps of the interval, *SIAM J. Appl. Math.* **35** (1978), no. 2, 260–267.
- [9] A. F. Ivanov and A. N. Sharkovsky, Oscillations in singularly perturbed delay equations, *Dynam. Rep. (New Series)* **1** (1992), 164–224.
- [10] A. J. Nicholson, An outline of the dynamics of animal populations, *Austral. J. Zool.* **2** (1954), 9–65.
- [11] W. S. C. Gurney, S. P. Blythe and R. M. Nisbet, Nicholson’s blowflies revisited, *Nature* **287** (1980), 17–21.
- [12] E. Braverman and D. Kinzbulatov, Nicholson’s blowflies equation with a distributed delay, *Can. Appl. Math. Quart.* **14** (2006), no. 2, 107–128.

- [13] J. Losson, M. C. Mackey, and A. Longtin, Solution multistability in first order nonlinear differential delay equations, *Chaos* **3** (1993), no. 2, 167–176.
- [14] E. Liz, E. Trofimchuk and S. Trofimchuk, Mackey-Glass type delay differential equations near the boundary of absolute stability, *J. Math. Anal. Appl.* **275** (2002), no. 2, 747–760.
- [15] E. Liz, C. Martínez and S. Trofimchuk, Attractivity properties of infinite delay Mackey-Glass type equations, *Differential Integral Equations* **15** (2002), no. 7, 875–896.
- [16] I. Kubiacyk and S. H. Saker, Oscillation and stability in nonlinear delay differential equations of population dynamics, *Math. Comput. Modelling* **35** (2002), no. 3-4, 295–301.
- [17] S. H. Saker, Oscillation and global attractivity in hematopoiesis model with delay time, *Appl. Math. Comput.* **136** (2003), no. 2-3, 241–250.
- [18] L. Berezansky and E. Braverman, Mackey-Glass equation with variable coefficients, *Comput. Math. Appl.* **51** (2006), no. 1, 1-16.
- [19] L. Berezansky and E. Braverman, On existence and attractivity of periodic solutions for the hematopoiesis equation, *Dyn. Contin. Discrete Impuls. Syst. Ser. A Math. Anal.* **13B** (2006), suppl., 103–116.
- [20] N. Bradul and L. Shaikhet, Stability of the positive point of equilibrium of Nicholson’s blowflies equation with stochastic perturbations: numerical analysis, *Discrete Dyn. Nat. Soc.* **2007**, Art. ID 92959, 25 pp.
- [21] C. Corduneanu, Functional Equations with Causal Operators. *Stability and Control: Theory, Methods and Applications*, **16**. Taylor & Francis, London, 2002.
- [22] I. Györi and G. Ladas, Oscillation Theory of Delay Differential Equations, Clarendon Press, Oxford, 1991.
- [23] J. Wei and M. Y. Li, Hopf bifurcation analysis in a delayed Nicholson blowflies equation, *Nonlinear Anal.* **60** (2005), no. 7, 1351–1367.
- [24] W. A. Coppel, The solution of equations by iteration, *Proc. Cambridge Philos. Soc.* **51** (1955), 41–43.
- [25] E. Liz, Local stability implies global stability in some one-dimensional discrete single-species models, *Discrete Contin. Dyn. Syst. Ser. B* **7** (2007), no. 1, 191–199.
- [26] L. Berezansky and E. Braverman, Linearized oscillation theory for a nonlinear equation with a distributed delay, *Math. Comput. Modelling* **48** (2008), no. 1-2, 287–304.